

**Children 0 months to 11 years**

**TYLENOL® Pediatric Products**

**DOSE: Every 4 hours as needed**

**Do NOT give more than 5 doses in 24 hours**

If possible, use weight to dose; otherwise, use age.

All Infants' TYLENOL® and Children's TYLENOL® products have the same strength of acetaminophen: **160 mg** (in each 5 mL, tablet, or powder).



**Infants' TYLENOL® Oral Suspension**

Active ingredient: acetaminophen 160 mg (in each 5 mL)

Available in: Grape Cherry Dye-Free Cherry

WEIGHT	6-11 lbs	12-17 lbs	18-23 lbs	24-35 lbs
AGE	0-3 months	4-11 months	12-23 months	2-3 years
DOSE	1.25 mL 	2.5 mL 	3.75 mL 	5 mL 

Use product only as directed.



**Children's TYLENOL® Chewables**

Active ingredient: acetaminophen 160 mg (in each chewable tablet)

Available in: Grape Bubblegum

WEIGHT	24-35 lbs	36-47 lbs	48-59 lbs	60-71 lbs	72-95 lbs
AGE	2-3 years	4-5 years	6-8 years	9-10 years	11 years
DOSE	1 tablet 	1½ tablets 	2 tablets 	2½ tablets 	3 tablets 

Use product only as directed.

mL = milliliter



**Children's TYLENOL® Oral Suspension**

Active ingredient: acetaminophen 160 mg (in each 5 mL)

Available in: Grape Cherry Dye-Free Cherry Bubblegum Strawberry

WEIGHT	24-35 lbs	36-47 lbs	48-59 lbs	60-71 lbs	72-95 lbs
AGE	2-3 years	4-5 years	6-8 years	9-10 years	11 years
DOSE	5 mL 	7.5 mL 	10 mL 	12.5 mL 	15 mL 

Use product only as directed.



**Children's TYLENOL® Dissolve Packs**

Active ingredient: acetaminophen 160 mg (in each powder)

Available in: Wild Berry

WEIGHT	under 48 lbs	48-59 lbs	60-71 lbs	72-95 lbs
AGE	under 6 years	6-8 years	9-10 years	11 years
DOSE	do not use	2 powders 	2 powders 	3 powders 

Use product only as directed.

**Important instructions for proper use:** Always read and follow the label on all TYLENOL® products. Use only the dosing device (dosing syringe or dosing cup) that came with the product. Do **NOT** use any other dosing device. Do **NOT** use with any other product containing acetaminophen.

**Adults and children 12 years and older**

**TYLENOL® Adult Products\***



**TYLENOL® Regular Strength Tablets and Liquid Gels**

Active ingredient: acetaminophen 325 mg (in each tablet or capsule)

Use products only as directed.

AGE	Children under 6 years	Children 6 years to under 12 years	Adults and children 12 years and over
DOSE AND FREQUENCY	Ask a doctor	1 tablet or capsule every 4-6 hours while symptoms last	2 tablets or capsules every 4-6 hours while symptoms last
MAXIMUM LABELED DOSE	Ask a doctor	Not to exceed 5 tablets or capsules in 24 hours	Not to exceed 10 tablets or capsules in 24 hours, unless directed by a doctor <i>Total labeled daily dose: 3250 mg</i>



**TYLENOL® Extra Strength Tablets and Dissolve Packs**

Active ingredient: acetaminophen 500 mg (in each tablet or powder)

Use products only as directed.

AGE	Adults and children 12 years and older
DOSE AND FREQUENCY	2 tablets or powders every 6 hours while symptoms last
MAXIMUM LABELED DOSE	Not to exceed 6 tablets or powders in 24 hours, unless directed by a doctor <i>Total labeled daily dose: 3000 mg</i>



**TYLENOL® 8HR Muscle Aches & Pain Tablets**

Active ingredient: acetaminophen 650 mg (in each tablet)†

Use product only as directed.

AGE	Adults and children 12 years and older
DOSE AND FREQUENCY‡	2 bi-layer tablets every 8 hours with water
MAXIMUM LABELED DOSE	Not to exceed 6 bi-layer tablets in 24 hours <i>Total labeled daily dose: 3900 mg</i>

**Important instructions for proper use:** Always read and follow the label on all TYLENOL® products. Do **NOT** use with any other product containing acetaminophen.

**Professional discretionary dosage:** If pain or fever persists at the total labeled daily dose, healthcare professionals may exercise their discretion and **recommend up to 4000 mg/day.**§

\*This is not a complete list of all available TYLENOL® adult products.

†Extended release.

‡Do not use in children under 12 years of age.

§The efficacy and safety of TYLENOL® at 4000 mg/day are well established.

Children 6 months to 11 years

**MOTRIN® Pediatric Products**

**DOSE: Every 6-8 hours as needed**

**Do NOT give more than 4 doses in 24 hours**

If possible, use weight to dose; otherwise, use age.



**Infants' MOTRIN® Concentrated Drops**

Active ingredient: ibuprofen (NSAID)\* 50 mg (in each 1.25 mL)

Available in: Berry Dye-Free Berry

WEIGHT	12-17 lbs	18-23 lbs
AGE	6-11 months	12-23 months
DOSE	1.25 mL 	1.875 mL 

Use product only as directed.



**Children's MOTRIN® Oral Suspension**

Active ingredient: ibuprofen (NSAID)\* 100 mg (in each 5 mL)

Available in: Grape Berry Dye-Free Berry Bubble Gum

WEIGHT	24-35 lbs	36-47 lbs	48-59 lbs	60-71 lbs	72-95 lbs
AGE	2-3 years	4-5 years	6-8 years	9-10 years	11 years
DOSE	5 mL 	7.5 mL 	10 mL 	12.5 mL 	15 mL 

Use product only as directed.

mL = milliliter



**Children's MOTRIN® Chewables**

Active ingredient: ibuprofen (NSAID)\* 100 mg (in each chewable tablet)

Available in: Grape Dye-Free Grape

WEIGHT	24-35 lbs	36-47 lbs	48-59 lbs	60-71 lbs	72-95 lbs
AGE	2-3 years	4-5 years	6-8 years	9-10 years	11 years
DOSE	1 tablet 	1½ tablets 	2 tablets 	2½ tablets 	3 tablets 

Use product only as directed.

**Important instructions for proper use:** Always read and follow the label on all MOTRIN® products. Use only the dosing device (dosing syringe or dosing cup) that came with the product. Do **NOT** use any other dosing device.

\*Nonsteroidal anti-inflammatory drug.

Adults and children 12 years and older

**MOTRIN® Adult Products\***



**MOTRIN® IB Liquid Gels**

Active ingredient: ibuprofen (NSAID)† 200 mg (in each capsule)

Use product only as directed.

AGE	Adults and children 12 years and older
DOSE AND FREQUENCY	1 capsule every 4-6 hours while symptoms last; if pain or fever does not respond to 1 capsule, 2 capsules may be used
MAXIMUM LABELED DOSE	Not to exceed 6 capsules in 24 hours, unless directed by a doctor <i>Total labeled daily dose: 1200 mg</i>



**MOTRIN® IB Caplets**

Active ingredient: ibuprofen (NSAID)† 200 mg (in each caplet)

Use product only as directed.

AGE	Adults and children 12 years and older
DOSE AND FREQUENCY	1 caplet every 4-6 hours while symptoms last; if pain or fever does not respond to 1 caplet, 2 caplets may be used
MAXIMUM LABELED DOSE	Not to exceed 6 caplets in 24 hours, unless directed by a doctor <i>Total labeled daily dose: 1200 mg</i>

**Important instructions for proper use:** Always read and follow the label on all MOTRIN® products.

†Nonsteroidal anti-inflammatory drug.

\*This is not a complete list of MOTRIN® adult products.

Find additional practice and patient resources at [TylenolProfessional.com](http://TylenolProfessional.com) and [JnJPediatrics.com](http://JnJPediatrics.com)